



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

particular cases of the following construction with ruler only, whenever a parallel to AB^1 is given: Join O , a point not on AB or the parallel, to A and to B meeting the parallel in D and C respectively. Let DB and CA meet in E , and OE meet AB in F . Join CF meeting DB in G , and let OG meet BA in H , etc. The points F, H , etc., are the same as those found before. This construction was given by Brianchon in 1818 in his *Application de la Théorie des Transversales*, page 37. He remarks: "Ce problème pourrait servir à se former, sur le terrain, une échelle de lever, si on n'avait pas à sa disposition une des mesures reçues, et qu'on connût d'ailleurs la longueur totale de la ligne prise pour échelle."

Lambert gave another construction in his *freye Perspective, oder Anweisung* . . . 1774, pages 173–174.

ARC.

PROBLEMS—SOLUTIONS

19 (Calculus) [1894, 165, 273–275]. Proposed by A. L. FOOTE, Merrick, N. Y.

A and B are in a circular room $2R = 30$ feet in diameter, A being at the center and B at the circumference. B runs around at the rate of $v = 600$ feet per minute and A pursues him at the rate of $u = 100$ feet per minute. How long will the race last, and how far will each have traveled till B is caught?

160 (Calculus) [1902, 271; 1903, 104–106]. Proposed by B. F. FINKEL, Drury College.

A dog at the vertex of a right conical hill pursues a fox at the foot of the hill. How far will the dog run to catch the fox, if the dog runs directly toward the fox at all times and the fox is continually running around the hill at its foot, the velocity of the dog being 6 feet per second, the velocity of the fox being 5 feet per second, the hill being 100 feet high and 200 feet in diameter at the base?

273 (Calculus) [1909, 76, 123–124; 1910, 221]. Proposed by J. SCHEFFER, Hagerstown, Md.

On one side of a circular pond a feet in radius is a duck. On the diametrically opposite side of the pond is a dog. Both swim at the same time, the duck swimming around the circumference of the pond at the rate of m feet a minute, the dog swimming directly towards the duck at the rate of n feet per minute. How far will the dog swim in overtaking the duck?

2801 [1920, 31]. Proposed by A. S. HATHAWAY, Houston, Texas.

A dog at the center of a circular pond makes straight for a duck which is swimming along the edge of the pond. If the rate of swimming of the dog is to the rate of swimming of the duck as $n : 1$, determine the equation of the curve of pursuit and the distance the dog swims to catch the duck.

I. REMARKS AND HISTORICAL NOTES BY R. C. ARCHIBALD AND H. P. MANNING, Brown University.

In 1732 Bourger read before the French Academy a memoir "Sur de nouvelles courbes ausquelles on peut donner le nom de Lignes de Poursuite"² in which he solved the following problem: "Trouver la courbe de poursuite, c'est-à-dire la courbe par laquelle un vaisseau doit en poursuivre un autre qui s'enfuit par une ligne droit, en supposant que les vitesses des deux vaisseaux soient toujours dans le même rapport."

Maupertuis gave³ a briefer solution of this and formulated also the following more general

¹ Or AA' bisected at B ; with this given Lambert showed, in 1774, that a parallel to AB can with ruler alone, readily be drawn through any point.

² *Histoire de l'Académie Royale des Sciences*, 1732. Paris, 1735, *Memoires*, pp. 1–14.

³ *Idem*, pp. 15–16.